

19



Europäisches Patentamt
European Patent Office
Office européen des brevets

11

Publication number:

**0 279 512
A2**

12

EUROPEAN PATENT APPLICATION

21

Application number: **88300376.6**

51

Int. Cl.4: **A61K 31/46** , **A61K 31/435** ,
A61K 31/445 , **A61K 31/55**

22

Date of filing: **18.01.88**

The title of the invention has been amended
(Guidelines for Examination in the EPO, A-III,
7.3).

30

Priority: **19.01.87 GB 8701022**

43

Date of publication of application:
24.08.88 Bulletin 88/34

64

Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

71

Applicant: **BEECHAM GROUP PLC**
Beecham House Great West Road
Brentford Middlesex TW8 9BD(GB)

72

Inventor: **Sanger, Gareth John**
Beecham Pharmaceuticals Coldharbour
Road
The Pinnacles Harlow Essex CM19 5AD(GB)
Inventor: **Marr, Helen Elizabeth**
Beecham Pharmaceuticals Coldharbour
Road
The Pinnacles Harlow Essex CM19 5AD(GB)

74

Representative: **Jones, Pauline et al**
Beecham Pharmaceuticals Patents & Trade
Marks Dept. Great Burgh Yew Tree Bottom
Road
Epsom Surrey KT18 5XQ(GB)

54

Use of certain 5-HT₃ receptor antagonists in the treatment of visceral pain.

57


A method of treatment of visceral pain in mammals, including humans, which method comprises the administration to the mammal in need of such treatment, an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof:

X-CO-Y-Z (I)


wherein

X is a group of formula (a), (b), (c), (d) or (e):


EP 0 279 512 A2




(a)




(b)



(c)



(d)



(e)

wherein

R_a to R_d are selected from hydrogen, halogen or hydroxy;


R_1 is hydrogen and R_2 is hydrogen or C_{1-4} alkyl; or

R_1 and R_2 together are a bond;


R_3 to R_7 are independently hydrogen or C_{1-6} alkyl; and

R_4 together with R_2 may be C_{2-7} polymethylene when R_1 is hydrogen;
either R_4 is C_{1-6} alkoxy;


R_8 is hydrogen;
 R_{10} is amino or C_{1-7} alkanoylamino; and
 R_{11} is halo or C_{1-6} alkylthio; or
 R_8 is hydrogen;
 R_9 is halo, C_{1-6} alkoxy or C_{1-6} alkyl;
 R_{10} is hydrogen or C_{1-6} alkoxy; and
 R_{11} is halo, C_{1-6} alkoxy or C_{1-6} alkyl;
 L is CH or N;
 Y is NH or O, with the proviso that Y is NH when X is (e) and R_8 is C_{1-6} alkoxy;
 Z is a group of formula (f), (g) or (h):



(f)



(g)



(h)

wherein
 n is 2 or 3;
 p and q are independently 1 to 3; and
 R_{12} or R_{13} is methyl or ethyl.

NOVEL TREATMENT


This invention relates to a method of treatment of visceral pain in mammals, including humans, and to the use of compounds in the preparation of a medicament for the treatment of visceral pain.

GB 2125398A, EP-A-200444, EP-A-247266, EP-A-235878, European Patent Application No. 87306548.6, EP-A-67770, EP-A-158265 and EP-A-158532 disclose classes of compounds which are 5-HT₂ receptor antagonists useful in the treatment of inter alia migraine, cluster headache, trigeminal neuralgia and emesis.

It has now been discovered that certain 5-HT₂ receptor antagonists, such as the above classes of compounds, are potentially useful in the treatment of visceral pain.


Visceral pain is caused by abnormal distension of hollow visceral organs. In particular, inflation of the colon of patients with irritable bowel syndrome will induce pain in various sites throughout the abdomen, mimicking the disease symptoms (Latimer et al, 1979, J. Behav. Med., 2, 285-295).

Accordingly, the present invention provides a method of treatment of visceral pain in mammals, including humans, which method comprises the administration to the mammal in need of such treatment, an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof:




wherein


X is a group of formula (a), (b), (c), (d) or (e):




(a)




(b)



(c)



(d)



(e)

wherein

R_a to R_d are selected from hydrogen, halogen or hydroxy;

R_1 is hydrogen and R_2 is hydrogen or C_{1-4} alkyl; or

R_1 and R_2 together are a bond;

R_3 to R_7 are independently hydrogen or C_{1-6} alkyl; and

R_4 together with R_2 may be C_{2-7} polymethylene when R_1 is hydrogen;

either R_6 is C_{1-8} alkoxy;

R_8 is hydrogen;

R_{10} is amino or C_{1-7} alkanoylamino; and

R_{11} is halo or C_{1-6} alkylthio; or

R_8 is hydrogen;

R_9 is halo, C_{1-6} alkoxy or C_{1-6} alkyl;


R_{10} is hydrogen or C_{1-6} alkoxy; and

5 R_{11} is halo, C_{1-6} alkoxy or C_{1-6} alkyl;


L is CH or N;

Y is NH or O, with the proviso that Y is NH when X is (e) and R_8 is C_{1-6} alkoxy;


Z is a group of formula (f), (g) or (h):



(f)



(g)



(h)

35 wherein

n is 2 or 3;

p and q are independently 1 to 3; and

R_{12} or R_{13} is methyl or ethyl.

Examples of moieties in alkyl or alkyl containing groups in R_1 to R_{11} include methyl, ethyl, n- and iso-propyl, n-, iso-, sec- and tert-butyl, preferably methyl.

40 Suitable examples of R_2 and R_4 when joined include C_2 , C_3 , C_4 , C_5 or C_6 polymethylene, preferably C_2 , C_3 , C_4 or C_5 polymethylene.

R_a to R_d are preferably selected from hydrogen, fluoro, chloro and hydroxy, most preferably hydrogen. R_b may be 5-, 6- or 7-chloro or fluoro.

45 When X is of sub-formula (a), R_1 and R_2 are preferably both hydrogen and one or both of R_2 and R_4 - (most preferably both) are alkyl groups, such as methyl, or are joined to form C_{2-7} polymethylene; or when one of R_2 and R_4 is hydrogen, the other is preferably ethyl or n- or iso-propyl.

When X is of sub-formula (b), R_5 is preferably hydrogen or a methyl or ethyl group.

When X is of sub-formula (c), one of CO-Y-Z and R_6 is attached at the 1-position and the other is attached at the 3-position as depicted in sub-formula (c), and R_6 is preferably methyl or ethyl.

50 When X is of sub-formula (d), R_7 is preferably methyl.

When X is of sub-formula (e), and R_8 is C_{1-6} alkoxy, R_8 is preferably methoxy, R_{10} is preferably amino and R_{11} is preferably chloro or bromo, most preferably chloro.

When X is of sub-formula (e), and R_8 is hydrogen, R_9 and R_{11} are preferably chloro or methyl and R_{10} is preferably hydrogen.

55 X is preferably a group of formula (b) and L is preferably N.

Y is preferably NH.

When Z is a group of sub-formula (f), n is 2 or 3, preferably 3 when X is of sub-formula (b) wherein L is

N.

When Z is a group of sub-formula (g) or (h), p and q are preferably 1 or 2.

The pharmaceutically acceptable salts of the compounds of the formula (I) include acid addition salts with conventional acids such as hydrochloric, hydrobromic, boric, phosphoric, sulphuric acids and pharmaceutically acceptable organic acids such as acetic, tartaric, lactic, maleic, citric, succinic, benzoic, ascorbic, methanesulphonic, α -keto glutaric, α -glycerophosphoric, and glucose-1-phosphoric acids.

The pharmaceutically acceptable salts of the compounds of the formula (I) are usually acid addition salts with acids such as hydrochloric, hydrobromic, phosphoric, sulphuric, citric, tartaric, lactic and acetic acid. Preferably the acid addition salt is the hydrochloride salt.

Pharmaceutically acceptable salts also include quaternary derivatives, examples of which include the compounds quaternised by compounds such as R_{20} -T wherein R_{20} is C_{1-8} alkyl, phenyl- C_{1-8} alkyl or C_5 - γ cycloalkyl, and T is a radical corresponding to an anion of an acid. Suitable examples of R_{20} include methyl, ethyl and n- and iso-propyl; and benzyl and phenethyl. Suitable examples of Z include halide such as chloride, bromide and iodide.

Pharmaceutically acceptable salts also include internal salts such as pharmaceutically acceptable N-oxides.

The compounds of the formula (I), and their pharmaceutically acceptable salts may also form pharmaceutically acceptable solvates, such as hydrates which are included wherever a compound of formula (I) or a salt thereof is herein referred to.

It will of course be realised that some of the compounds of the formula (I) have chiral or prochiral centres, and thus are capable of existing in a number of stereoisomeric forms, including enantiomers. The invention extends to each of these stereoisomeric forms (including enantiomers), and to mixtures thereof (including racemates). The different stereoisomeric forms may be separated one from the other by the usual methods.

Compounds of the formula (I) and their salts may be prepared in accordance with the methods described in the abovementioned UK and European Patent references, the subject matter of which are incorporated herein by reference.

The administration of the compound of formula (I) or a pharmaceutically acceptable salt thereof may be by way of oral or parenteral administration.

An amount effective to treat the disorders hereinbefore described depends on the relative efficacies of the compounds of the invention, the nature and severity of the disorder being treated and the weight of the mammal. However, a unit dose for a 70kg adult will normally contain 0.1 to 100mg for example 0.2 to 50mg, of the compound of formula (I) or a pharmaceutically acceptable salt thereof. Unit doses may be administered once or more than once a day, for example, 2, 3 or 4 times a day, more usually 1 to 3 times a day, that is in the range of approximately 0.0002 to 5 mg/kg/day, more usually 0.0004 to 2.5 mg/kg/day.

No adverse toxicological effects are indicated at any of the aforementioned dosage ranges.

It is preferred that the compound of formula (I) or a pharmaceutically acceptable salt thereof is administered in the form of a unit dose pharmaceutical composition in which is combined with a pharmaceutically acceptable carrier.

Such compositions are prepared by admixture and are suitably adapted for oral or parenteral administration, and as such may be in the form of tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutable powders, injectable and infusible solutions or suspensions or suppositories. Orally administrable compositions are preferred, since they are more convenient for general use.

Tablets and capsules for oral administration are usually presented in a unit dose, and contain conventional excipients such as binding agents, fillers, diluents, tableting agents, lubricants, disintegrants, colourants, flavourings, and wetting agents. The tablets may be coated according to well-known methods in the art, for example with an enteric coating.

Suitable fillers for use include cellulose, mannitol, lactose and other similar agents. Suitable disintegrants include starch, polyvinylpyrrolidone and starch derivatives such as sodium starch glycolate.

Suitable lubricants include, for example, magnesium stearate.

Suitable pharmaceutically acceptable wetting agents include sodium lauryl sulphate. Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, gelatin, hydroxyethylcellulose, carboxymethylcellulose, aluminium stearate gel or hydrogenated edible fats, emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example, almond oil, fractionated coconut oil, oily esters such as esters of glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or

propyl p-hydroxybenzoate or sorbic acid, and if desired conventional flavouring or colouring agents.

Oral liquid preparations are usually in the form of aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs or are presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, emulsifying agents, non-aqueous vehicles (which may include edible oils), preservatives, and flavouring or colouring agents.

The oral compositions may be prepared by conventional methods of blending, filling or tableting. Repeated blending operations may be used to distribute the active agent throughout those compositions employing large quantities of fillers. Such operations are, of course, conventional in the art.

For parenteral administration, fluid unit dose forms are prepared containing a compound of the present invention and a sterile vehicle. The compound, depending on the vehicle and the concentration, can be either suspended or dissolved. Parenteral solutions vehicle and filter sterilising before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering agents are also dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum.

Parenteral suspensions are prepared in substantially the same manner except that the compound is suspended in the vehicle instead of being dissolved and sterilised by exposure of ethylene oxide before suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound.

The present invention also provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof in the preparation of a medicament for use in the treatment of visceral pain in mammals, including humans. Such treatment may be carried out in the manner as hereinbefore described.

The present invention further provides a pharmaceutical composition for use in the treatment of visceral pain which comprises an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier. Such compositions may be prepared in the manner as hereinbefore described.

The following pharmacological data illustrate the invention.

Compound E6 is the Compound of Example 6 of EP-A-200444, N-(endo-9-methyl-9-azabicyclo[3.3.1]-non-3-yl)-1-methyl-indazole-3-carboxamide monohydrochloride (also known as BRL 43694A).

ICS 205-930 is the Compound of Example A-5 of GB 2125398A, 2-methoxy-indol-3-yl carboxylic acid (endo-8-methyl-8-azabicyclo[3.2.1]oct-3-yl)ester.

Compound E5 is the Compound of Example 5 of EP-A-247266, endo-N-(8-methyl-8-azabicyclo[3.2.1]-oct-3-yl)-2,3-dihydro-3,3-dimethylindole-1-carboxamide.

Pharmacological Data

Visceral pain in man or in animals is associated with a range of pseudoaffective, pain-related responses which are easily monitored. For example, distension of anaesthetised rat ileum induces a fall in blood pressure which can be blocked by morphine or by capsaicin, indicating that it is part of the pain reflex (Lembeck and Skofitsch, 1982, Naunyn-Schmiedeberg's Arch. Pharmac., 321, 179-183). Furthermore, nociceptive stimuli directed to the intestine can reflexly suppress gastric motility (Abrahamsson et al, 1979, Scand. J. Gastroenterol., 14, 101-106). Compounds which selectively block such pseudoaffective responses may therefore prevent both visceral pain (e.g. pain associated with the irritable bowel syndrome, gall stones, kidney stones, etc) and the paralytic ileus (suppression of gastrointestinal motility) associated with abdominal surgery.

Method

Rats were prepared under urethane anaesthesia for measurement of blood pressure via a cannula in the carotid artery. The whole stomach was intubated with a cannula down the oesophagus and ligated distal to the pyloric sphincter before being filled with 5ml saline. A 3-4cm segment of the duodenum was isolated immediately below the stomach. This segment was connected orally to saline-filled pressure reservoirs with drainage occurring aborally. Distension of the duodenum with a pressure of 75cmH₂O induced a fall in blood

pressure and a fall in intragastric pressure, which could be repeated over a series of distensions. Intravenous (i.v.) injection of E6 $10\mu\text{g kg}^{-1}$ reduced both the fall in blood pressure and intragastric pressure caused by duodenal distension.

The results are shown in Tables 1 to 3.

TABLE 1

Inhibition by E6 of the fall in blood pressure and intragastric pressure induced by duodenal distension in the anaesthetised rat

Time (min) after administration of E6 $1\mu\text{g/kg}$ (n=6)	% Inhibition of fall in blood pressure	% Inhibition of fall in intragastric pressure
5	34 ± 16	19 ± 8
10	$37 \pm 14^*$	0 ± 13
15	32 ± 14	10 ± 11
20	35 ± 16	9 ± 9

Time (min) after administration of E6 $10\mu\text{g/kg}$ (n=9)	% Inhibition of fall in blood pressure	% Inhibition of fall in intragastric pressure
5	2 ± 17	23 ± 10
10	$37 \pm 8^{***}$	$49 \pm 15^{**}$
15	$47 \pm 14^{**}$	$66 \pm 15^{***}$
20	$56 \pm 8^{***}$	$74 \pm 14^{***}$

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.005$
Student's 't' test

TABLE 2

Inhibition by ICS 205-930 of the fall in blood pressure and intragastric pressure induced by duodenal distension in the anaesthetised rat

Time (min) after administration of ICS 205-930 100µg/kg	% Inhibition of fall in blood pressure (n=5)	% Inhibition of fall in intragastric pressure (n=4)
5	8 ± 27	5 ± 13
10	33 ± 7**	37 ± 6**
15	47 ± 18	46 ± 8**
20	45 ± 16*	27 ± 20

*P<0.05; **P<0.01
Student's 't' test

TABLE 3

Inhibition by E5 of the fall in blood pressure and intragastric pressure induced by duodenal distension in the anaesthetised rat

Time (min) after administration of E5 10µg/kg	% Inhibition of fall in blood pressure (n=9)	% Inhibition of fall in intragastric pressure (n=8)
5	36 ± 15*	16 ± 5
10	41 ± 18	63 ± 14***
15	37 ± 22	51 ± 14**
20	32 ± 19	47 ± 17*

*P<0.05; **P<0.01; ***P<0.005
Student's 't' test

Claims

1. Use of a compound of formula (I) or a pharmaceutically acceptable salt thereof:

X-CO-Y-Z (I)


wherein

X is a group of formula (a), (b), (c), (d) or (e):

5

10


15



(a)

20


25



(b)

30


35



(c)

40


45



(d)

50

55



(e)

wherein

R_a to R_d are selected from hydrogen, halogen or hydroxy;

R₁ is hydrogen and R₂ is hydrogen or C₁₋₄ alkyl; or

R_1 and R_2 together are a bond;

R_3 to R_7 are independently hydrogen or C_{1-6} alkyl; and

R_4 together with R_2 may be C_{2-7} polymethylene when R_1 is hydrogen;

either R_8 is C_{1-6} alkoxy;

R_9 is hydrogen;

R_{10} is amino or C_{1-7} alkanoylamino; and

R_{11} is halo or C_{1-6} alkylthio; or

R_8 is hydrogen;

R_9 is halo, C_{1-6} alkoxy or C_{1-6} alkyl;


R_{10} is hydrogen or C_{1-6} alkoxy; and

R_{11} is halo, C_{1-6} alkoxy or C_{1-6} alkyl;


L is CH or N;

Y is NH or O, with the proviso that Y is NH when X is (e) and R_8 is C_{1-6} alkoxy;


Z is a group of formula (f), (g) or (h):



(f)



(g)



(h)

wherein

n is 2 or 3;

p and q are independently 1 to 3; and

R_{12} or R_{13} is methyl or ethyl; in the manufacture of a medicament for the treatment of visceral pain.

2. A use according to claim 1 wherein X is of sub-formula (a) as defined in claim 1, and R_1 and R_2 are both hydrogen and either R_3 and R_4 are both methyl groups or are joined to form C_{2-7} polymethylene, or one of R_3 and R_4 is hydrogen and the other is ethyl or *n*- or *iso*-propyl.

3. A use according to claim 1 wherein X is of sub-formula (b) as defined in claim 1 and R_5 is hydrogen or a methyl or ethyl group.

4. A use according to claim 1 wherein Z is of sub-formula (f) as defined in claim 1, and n is 2 or 3.

5. A use according to claim 1 wherein the compound of formula (I) is N-(endo-9-methyl-9-azabicyclo-[3.3.1]non-3-yl)-1-methyl-indazole-3-carboxamide or a pharmaceutically acceptable salt thereof.

6. A use according to claim 1 wherein the compound of formula (I) is 2-methoxy-indole-3-yl carboxylic acid (endo-8-methyl-8-azabicyclo[3.2.1]oct-3-yl)ester, or a pharmaceutically acceptable salt thereof.

7. A use according to claim 1 wherein the compound of formula (I) is endo-N-(8-methyl-8-azabicyclo-[3.2.1]oct-3-yl)-2,3-dihydro-3,3-dimethylindole-1-carboxamide.



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number: **0 279 512 A3**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **88300376.6**

(51) Int. Cl.⁵: **A61K 31/46, A61K 31/435,
A61K 31/445, A61K 31/55**

(22) Date of filing: **18.01.88**

(30) Priority: **19.01.87 GB 8701022**

(43) Date of publication of application:
24.08.88 Bulletin 88/34

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(88) Date of deferred publication of the search report:
16.09.92 Bulletin 92/38


(71) Applicant: **BEECHAM GROUP PLC**
Beecham House Great West Road
Brentford Middlesex TW8 9BD(GB)

(72) Inventor: **Sanger, Gareth John**
Beecham Pharmaceuticals Coldharbour
Road
The Pinnacles Harlow Essex CM19 5AD(GB)
Inventor: **Marr, Helen Elizabeth**
Beecham Pharmaceuticals Coldharbour
Road
The Pinnacles Harlow Essex CM19 5AD(GB)

(74) Representative: **Jones, Pauline et al**
Beecham Pharmaceuticals Patents & Trade
Marks Dept. Great Burgh Yew Tree Bottom
Road
Epsom Surrey KT18 5XQ(GB)

(54) **Use of certain 5-HT₃ receptor antagonists in the treatment of visceral pain.**


(57) A method of treatment of visceral pain in mammals, including humans, which method comprises the administration to the mammal in need of such treatment, an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof:




wherein

X is a group of formula (a), (b), (c), (d) or (e):


EP 0 279 512 A3




(a)




(b)



(c)



(d)



(e)

wherein


R_a to R_d are selected from hydrogen, halogen or hydroxy;

R_1 is hydrogen and R_2 is hydrogen or C_{1-4} alkyl; or


R_1 and R_2 together are a bond;

R_3 to R_7 are independently hydrogen or C_{1-6} alkyl; and


R_4 together with R_2 may be C_{2-7} polymethylene when R_1 is hydrogen;
 either R_8 is C_{1-6} alkoxy;
 R_9 is hydrogen;
 R_{10} is amino or C_{1-7} alkanoylamino; and
 R_{11} is halo or C_{1-6} alkylthio; or
 R_8 is hydrogen;
 R_9 is halo, C_{1-6} alkoxy or C_{1-6} alkyl;
 R_{10} is hydrogen or C_{1-6} alkoxy; and
 R_{11} is halo, C_{1-6} alkoxy or C_{1-6} alkyl;
 L is CH or N;
 Y is NH or O, with the proviso that Y is NH when X is (e) and R_8 is C_{1-6} alkoxy;
 Z is a group of formula (f), (g) or (h):



(f)



(g)



(h)

wherein

n is 2 or 3;

p and q are independently 1 to 3; and

R_{12} or R_{13} is methyl or ethyl.



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention
shall be considered, for the purposes of subsequent
proceedings, as the European search report

EP 88 30 0376

DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
Y	NATURE, vol. 316, no. 6024, July 11-17, 1985, pages 126-131; B.P. RICHARDSON et al.: "Identification of serotonin M-receptor subtypes and their specific blockade by a new class of drugs" * Summary * ---	1-7	A 61 K 31/46 A 61 K 31/435 A 61 K 31/445 A 61 K 31/55
Y	WO-A-8 502 847 (SANDOZ AG) * Page 30, line 4; claim 8 * ---	1-7	
Y,D	GB-A-2 125 398 (SANDOZ LTD) * Page 13, line 35 * ---	1-7	
Y,D	EP-A-0 067 770 (MERRELL TORAUDE ET COMPAGNIE) * Page 8, lines 22-25 * ---	1-7	
Y	EP-A-0 220 011 (BEECHAM GROUP PLC) * Page 2, lines 10-12 * ---	1-7	
Y	EP-A-0 189 002 (SANDOZ AG) * Pages 18-19 * --- -/-	1-7	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			A 61 K 31/00

INCOMPLETE SEARCH

The Search Division considers that the present European patent application does not comply with
the provisions of the European Patent Convention to such an extent that it is not possible to carry
out a meaningful search into the state of the art on the basis of some of the claims

Claims searched completely :

Claims searched incompletely :

Claims not searched :

Reason for the limitation of the search:

Place of search

THE HAGUE

Date of completion of the search

16-06-1992

Examiner

THEUNS H.G.

CATEGORY OF CITED DOCUMENTS

X : particularly relevant if taken alone
Y : particularly relevant if combined with another
document of the same category
A : technological background
O : non-written disclosure
P : intermediate document

T : theory or principle underlying the invention
E : earlier patent document, but published on, or
after the filing date
D : document cited in the application
L : document cited for other reasons

& : member of the same patent family, corresponding
document



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Page 2

Application Number

EP 88 30 0376

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
Y	EP-A-0 201 165 (BEECHAM GROUP PLC) * Page 1, lines 28-29 * ---	1-7	
Y,D	EP-A-0 200 444 (BEECHAM GROUP PLC) * Page 21, lines 9-10 * ---	1-7	
Y	EP-A-0 223 385 (BEECHAM GROUP PLC) * Page 7, lines 37-3 * ---	1-7	
A	THE MERCK MANUAL OF DIAGNOSIS AND THERAPY, 14th Ed., 1982, Merck & Co., Inc., Rahway, N.J., US * Pages 736-73, 790-791 and 1293 * -----	1-7	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)

EPO FORM 1503 03.82 (P0410)